

## SECTION I: SCENARIO OVERVIEW

<b>Scenario Title:</b>	Adult Med-Surg: Blood Transfusion Reaction-Anaphylaxis	
Original Scenario Developer(s):	Sally Scofield, MSN, RN <a href="mailto:sscofield@ohlone.edu">sscofield@ohlone.edu</a> Kathy Dewan, MSN, RN <a href="mailto:kdewan@ohlone.edu">kdewan@ohlone.edu</a> Gale Carli, EdD, RN <a href="mailto:gcarli@ohlone.edu">gcarli@ohlone.edu</a>	
Date - original scenario	03/07	
Validation:	07/08	
Revision Dates:	08/08, 10/12, 07/14, 3/18 (mmiller@nurse-edconsulting.com)	
Pilot testing:	07/08	
QSEN revision:	10/12 Barbara Durham, MSN, RN, CNE <a href="mailto:bdurham@hartnell.edu">bdurham@hartnell.edu</a> Marjorie Miller, MA, RN, CHSE <a href="mailto:mmiller@nurse-edconsulting.com">mmiller@nurse-edconsulting.com</a>	
<b>Estimated Scenario Time:</b> 15 minutes <b>Debriefing time:</b> 30 minutes		
<p><u>Target group:</u> Pre-licensure nursing students, new graduates. Key learning involves introducing students to monitoring priorities, recognizing &amp; responding to abnormal findings &amp; documentation involved with blood administration and potential transfusion reaction.</p> <p><u>Core case:</u> Blood transfusion reaction-anaphylaxis</p> <p><u>QSEN Competencies:</u> Safety, Teamwork and Collaboration, Evidence-based practice, Patient-centered care</p> <p><u>Brief Summary of Case:</u> James Patterson, a 55-year old man with newly diagnosed non-small cell cancer of the lung, is admitted with complaints of fever, weakness and sore throat for two days. He informs the nurse that he has felt very tired, bruises easily and has no appetite because of sore throat since his first chemotherapy 1 ½ weeks ago. Mr. Patterson describes getting weaker every day and slightly short of breath with activity. CBC shows pancytopenia. MD orders to transfuse 2 units of PRBC. This is Mr. Patterson’s first blood transfusion. Learners are expected to safely initiate a blood transfusion according to protocol, monitor vital signs, recognize acute hemolytic reaction, stop the blood, call for help and notify physician utilizing SBAR communication. Further the learner should take and validate orders via EMR and administer stat medications.</p>		

EVIDENCE BASE / REFERENCES (APA Format)
Cronenwett, L, Sherwood, G., Barnsteiner, J., et al. (2007). Quality and safety education for nurses. <i>Nursing Outlook</i> , 55(3), 122-131. doi:10.1016/j.outlook.2007.02.o06
Pham, Huy Phu (2017-revised) Allergic and anaphylactic transfusion reaction. Pathology Outlines.com website. <a href="http://www.pathologyoutlines.com/topic/transfusionmedallergic.html">http://www.pathologyoutlines.com/topic/transfusionmedallergic.html</a> Accessed March 8, 2018.
Rome, S.I. (2016). Hematologic problems. In S.L. Lewis, M. Heitkemper, S. R. Dirksen, P. G. O’Brien & L. Bucher (Eds.), <i>Medical-Surgical Nursing: Assessment and management of clinical problems</i> (pp. 732-734). St. Louis, MO: Elsevier.
Sandler, S. G. (2017). Transfusion reactions. Treatment & Management. Retrieved from <a href="http://emedicine.medscape.com">http://emedicine.medscape.com</a>

## SECTION II: CURRICULUM INTEGRATION

### A. SCENARIO LEARNING OBJECTIVES

#### Learning Outcomes

1. Integrate understanding of multiple dimensions of patient centered care
2. Assess own level of communication skill in encounters with patients and families.
3. Apply clinical decision making skills in interpreting and analyzing data in evolving situations.
4. Prioritize interventions based on accurate interpretation of assessment data and initiates request for help when appropriate to situation.

#### Specific Learning Objectives

1. Confirms physician orders for blood transfusion and informed consents are in the chart
2. Gathers relevant patient and contextual data to identify patient’s current problem.
3. Apply principles of hand hygiene, infection control and personal protection during the initiation of a blood transfusion.
4. Demonstrates situational awareness and recognizes acute changes that warrant immediate intervention.
5. Prioritizes and implements interventions in response to transfusion reaction according to institution-specific policies.
6. Assessment and management of a blood transfusion reaction, including documentation.
7. Provides patient-centered care (teaching/learning opportunities, therapeutic communication) and reassures patient throughout simulation with clear, calm statements of action.
8. Communicates effectively with nursing and members of inter-professional team regarding the management of a patient with a transfusion reaction.

#### Critical Learner Actions

1. Perform hand hygiene, don gloves, and introduces self / identify patient with 2 patient identifiers.
2. Verifies completion of required documents prior to initiating transfusion (consent, availability of blood).
3. Confirms patient understanding of procedure.
4. Follows institution-specific policies for initiation of a blood transfusion.
5. Assessment and documentation of vital signs, oxygen saturation, s/s of blood transfusion reaction.
6. Recognize signs and symptoms and identifies transfusion reaction.
7. Immediate response and management of transfusion reaction according to institution specific policy.
8. Use standardized communication tool (i.e. SBAR) to communicate patient status to interprofessional team.

### B. PRE-SCENARIO LEARNER ACTIVITIES

#### Prerequisite Competencies

Required prior to participating in the scenario

Knowledge	Skills/ Attitudes
<input type="checkbox"/> Focused assessment (Adult) including patient understanding of the procedure.	<input type="checkbox"/> Signs and symptoms indicating the need for blood transfusion.
<input type="checkbox"/> IV infusion/set up of blood-Y tubing and administration of blood products <input type="checkbox"/> Administration of IV medications	<input type="checkbox"/> Knowledge of policies and procedures / documentation required for blood product administration.
<input type="checkbox"/> Knowledge of policies and procedures for blood product administration.	<input type="checkbox"/> RN responsibilities and rationale before, during and after blood transfusion.
<input type="checkbox"/> Universal precautions	<input type="checkbox"/> Signs and symptoms of a blood transfusion reaction.

### SECTION III: SCENARIO SCRIPT

**A. Case summary**

James Patterson, a 55-year old man with newly diagnosed non-small cell cancer of the lung, is admitted with complaints of fever, weakness and sore throat for two days. He informs the nurse that he bruises very easily. He also informs the nurse that he completed his first round of chemotherapy about a week and a half ago. Since then he has felt very tired and has no appetite. He could barely eat because of sore throat. Mr. Patterson describes getting weaker every day and slightly short of breath with activity. CBC shows pancytopenia. MD orders to transfuse 2 units of PRBC. This is Mr. Patterson’s first blood transfusion.

**B. Key contextual details**

The scenario takes place mid-shift on an oncology floor. The unit is fully staffed. Blood bank called indicating that the blood for Mr. Patterson is available to be picked up. A charge nurse and resource nurse are available.

**C. Scenario Cast**

Patient/ Client	<input type="checkbox"/> High fidelity simulator	
	<input type="checkbox"/> Mid-level simulator	
	<input type="checkbox"/> Task trainer	
	<input type="checkbox"/> Hybrid (Blended simulator)	
	<input type="checkbox"/> Standardized patient	
Role	Brief Descriptor (Optional)	Standardized Participant (SP) or Learner (L)
RN 1		L
RN 2		L
Charge nurse		SP
Physician	Will answer phone call and provide orders for treatment of reaction and/or remind nurse of policy for transfusion reaction. See case flow for script	SP
Family member	Add for complexity if needed	SP

D. Patient/Client Profile				
Last name:	Patterson		First name:	James
Gender: M	Age: 50	Ht: 6'0"	Wt: 187 lbs (85kg)	Code Status: Full
Spiritual Practice: None		Ethnicity: Caucasian		Primary Language spoken: English
1. History of present illness				
Newly diagnosed with non-small cell cancer of the lung. Complains of fever, weakness, and sore throat for two days. He has multiple ecchymotic areas on his arms and legs. He c/o feeling tired and has lack of appetite. He completed his first round of chemotherapy 1 ½ weeks ago. CBC shows pancytopenia.				
<b>Primary Medical Diagnosis</b>		Lung Cancer, Pancytopenia following chemotherapy		

2. Review of Systems	
CNS	Awake, alert, oriented
Cardiovascular	S1/S2
Pulmonary	Diminished breath sounds, bases bilaterally, A and P
Renal/Hepatic	Urine yellow, concentrated
Gastrointestinal	Mild nausea, positive bowel sounds
Endocrine	No problems
Heme/Coag	Pancytopenic
Musculoskeletal	Equal bilateral
Integument	Generalized petechiae, hematoma in upper arms, pale skin and sclerae
Developmental Hx	Adult milestones obtained
Psychiatric Hx	None contributory
Social Hx	Smoked until time of diagnosis, 100 pack-year smoking history
Alternative/ Complementary Medicine Hx	None

Medication allergies:	NKDA	Reaction:	
Food/other allergies:	NKFA	Reaction:	

3. Current medication	Drug	Dose	Route	Frequency
	Allopurinol	300 mg	oral	daily
	Procrit	100 mg	Sub-q	Monday / Wednesday
	Tylenol	325 mg 1-2 tabs	oral	Q 6 h prn fever or HA

4. Laboratory, Diagnostic Study Results					
Na: 134	K: 4.0	Cl: 100	HCO3: 26	BUN: 28	Cr: 0.9
Ca:	Mg:	Phos:	Glucose:	HgA1C:	
Hgb: 7.4 gm/dL	Hct: 22.2 %	Plt: 90,000	WBC: 2.5mm <sup>3</sup>	ABO Blood Type:	
PT	PTT	INR	Troponin:	BNP:	
Ammonia:	Amylase:	Lipase:	Albumin:	Lactate:	
ABG-pH:	paO2:	paCO2:	HCO3/BE:	SaO2:	
VDRL:	GBS:	Herpes:	HIV:		
CXR:	ECG:				
CT:	MRI:				
Other: ABO blood type: A positive					

E. Baseline Simulator/Standardized Patient State (This may vary from the baseline data provided to learners)					
1. Initial physical appearance					
Gender: M		Attire: patient gown			
Alterations in appearance (moulage): Pale color; multiple small bruised areas to arms / legs if possible					
X	ID band present, accurate information		ID band present, inaccurate information		ID band absent or not applicable
	Allergy band present, accurate information		Allergy band present, inaccurate information	X	Allergy band absent or not applicable

2. Initial Vital Signs Monitor display in simulation action room:					
	No monitor display	X	Monitor on, but no data displayed		Monitor on, standard display

BP: 110/64	HR: 100	RR: 24	T: 99.8 °F	SpO <sup>2</sup> : 93%	
CVP:	PAS:	PAD:	PCWP:	CO:	
AIRWAY:	ETCO <sup>2</sup> :	FHR:			
Lungs: Sounds/mechanics	Left: CLEAR but diminished		Right: CLEAR but diminished		
Heart:	Sounds:	S1/S2			
	ECG rhythm:	Sinus rhythm/sinus tachycardia			
	Other:				
Bowel sounds:	Present		Other:		

3. Initial Intravenous line set up						
	Saline lock #1	Site:				IV patent (Y/N)
X	IV #1	Site: RA	20 G	Fluid type: D5½NS	Initial rate: 100 ml/hr	X IV patent (Y/N)
X	Main					
	Piggyback					
	IV #2	Site:		Fluid type:	Initial rate:	IV patent (Y/N)
	Main					
	Piggyback					
4. Initial Non-invasive monitors set up						
X	NIBP	X	ECG First lead:		ECG Second lead:	
X	Pulse oximeter		Temp monitor/type		Other:	
5. Initial Hemodynamic monitors set up						
	A-line Site:		Catheter/tubing Patency (Y/N)	CVP Site:	PAC Site:	
6. Other monitors/devices						
	Foley catheter	Amount:	Appearance of urine:			
	Epidural catheter	Infusion pump:	Pump settings:			
	Fetal Heart rate monitor/tocometer	Internal	External			

<b>Environment, Equipment, Essential props</b>	
Recommend standardized set ups for each commonly simulated environment	
<b>1. Scenario setting: (example: patient room, home, ED, lobby)</b>	
Med/Surg / Telemetry unit, patient room	
Simulated blood transfusion 1 unit	

<b>2. Equipment, supplies, monitors</b>				
(In simulation action room or available in adjacent core storage rooms)				
X	Bedpan/ Urinal	Foley catheter kit	Straight cath. kit	Incentive spirometer
X	IV Infusion pump	Feeding pump	Pressure bag	Wall suction
	Nasogastric tube	ETT suction catheters	Oral suction catheters	Chest tube insertion kit
	Defibrillator	Code Cart	12-lead ECG	Chest tube equip
	PCA infusion pump	Epidural infusion pump	Central line Insertion Kit	Dressing Δ equipment
X	IV fluid	IV fluid additives:	Blood "Y" set for transfusion	X Blood product ABO Type: A+ # of units: 1
X	Type: D5½NS NS for blood			

<b>3. Respiratory therapy equipment/devices</b>					
X	Nasal cannula	Face tent	X	Simple Face Mask	Non re-breather mask
X	BVM/Ambu bag	Nebulizer tx kit		Flowmeters (extra supply)	

<b>4. Documentation and Order Forms</b>						
X	Health Care Provider orders	X	Med Admin Record	H & P	X	Lab Results
	Progress Notes	X	Graphic record	Anesthesia/PACU record		ED Record
	Medication reconciliation		Transfer orders	X	Standing (protocol) orders	ICU flow sheet
X	Nurses' Notes		Dx test reports		Code Record	Prenatal record
X	Actual medical record binder, constructed per institutional guidelines or EMR			Other	Describe: Transfusion Record	

<b>5. Medications (to be available in sim action room)</b>								
#	Medication	Dosage	Route		#	Medication	Dosage	Route
	Benadryl	50 mg	IVP			NS bolus	500ml	IV
	Epinephrine	0.5 ml of 1:1000	SQ	OR		Epinephrine	0.5 ml of 1:10,000	IV

**CASE FLOW / TRIGGERS/ SCENARIO DEVELOPMENT STATES**

**Initiation of Scenario :** Report is given to the nurse by the off-going shift (who had to leave mid shift due to a family emergency):

Mr. Patterson has non-small cell cancer of the lung. He completed his first round of chemotherapy 1 ½ weeks ago. He states having a fever, weakness, and sore throat for two days. He is tired and has no appetite. Has shortness of breath on activity. Patient seen in ED last night and admitted for further workup. D<sub>5</sub> ½ NS at 100 ml/hr started in ED 18g RA. Labs drawn in ED: CBC, Chem panel, electrolytes, BC x2, Type and cross match 2 units PRBC. His H/H were low (7.4 gm/dL and 22.2%). The MD ordered to transfuse 2 units PRBC's. His vital signs have been stable and he voided 100 cc amber urine after he was admitted. "I called for the first unit. You will need to start his transfusion."

STATE / PATIENT STATUS	DESIRED LEARNER ACTIONS & TRIGGERS TO MOVE TO NEXT STATE		
<p><b>1. Baseline</b></p> <p>Bruising on extremities Pale Dry mucous membranes</p> <p>Responsive but fatigued Flat vocal tone</p>	<p><b>Operator</b></p> <p>HR: 104, pulse weak RR: 20, diminished at bases, BP: 110/64 T: 99.2° F O<sub>2</sub> Sat: 93% on 2L NC Occasional dry cough</p> <p><b>Triggers:</b> Complete focused assessments in 2 to 3 min</p>	<p><b>Learner Actions</b></p> <ul style="list-style-type: none"> <li>• Washes hands/ID patient/ introduce self</li> <li>• Obtains/ documents baseline VS/ O<sub>2</sub> sats</li> <li>• Performs a focused assessment (LOC, breath sounds, SOB and other symptoms indicating need for transfusion)</li> <li>• Assesses patient's understanding of procedure</li> </ul>	<p><b>Debriefing Points:</b></p> <ul style="list-style-type: none"> <li>• Strategies for adhering to NPSG regarding patient identification.</li> <li>• Initial focused assessment compared to what was reported</li> <li>• Importance of obtaining baseline vital signs and the relevance of the values               <ul style="list-style-type: none"> <li>○ Anemia causes SOB, dec O<sub>2</sub> sats, increased workload on heart (tachycardia), dizziness, weakness, fatigue etc.</li> </ul> </li> <li>• Principles of patient centered care: asking the patient to "teach back".</li> </ul>



STATE / PATIENT STATUS	DESIRED ACTIONS & TRIGGERS TO MOVE TO NEXT STATE		
<p>2.</p> <p>Patient status unchanged</p> <p>Patient responds lethargically asks about transfusion -- "why do I need that?"</p>	<p><b>Operator:</b></p> <p>VS unchanged</p> <p><b>Triggers:</b></p> <p>Begins transfusion within 5 min after receiving unit</p>	<p><b>Learner Actions:</b></p> <ul style="list-style-type: none"> <li>• Reviews MD order for blood transfusion</li> <li>• Checks chart for required documentation for blood administration (consent, refusal, religious preferences, ABO compatibility, labs)</li> <li>• Inform patient of procedure (purpose, duration), explains assessment findings to report</li> <li>• Sends a helper to pick up blood while preparing IV tubing for blood transfusion (holds maintenance IV as noted in chart)</li> <li>• Checks blood at bedside with another nurse</li> <li>• Begin transfusion</li> </ul>	<p><b>Debriefing Points:</b></p> <ul style="list-style-type: none"> <li>• Institution specific policy/procedure must be followed for blood administration.</li> <li>• Key teaching points prior to administering blood (i.e. signs and symptoms to report)</li> <li>• Need for safety checks, two patient identifiers according to NPSG.</li> <li>• Describe the decision-making process and priority setting</li> <li>• Correct drug/dose calculation               <ul style="list-style-type: none"> <li>○ initial rate versus maintenance rate</li> </ul> </li> </ul>

STATE / PATIENT STATUS	DESIRED ACTIONS & TRIGGERS TO MOVE TO NEXT STATE		
<p><b>3.</b></p> <p><b>10-15 minute time period elapses (condensed over a 5 minute period)</b></p> <p>Moaning, gasping with interrupted speech and shortness of breath</p> <p><b>Progressively states:</b>            “My IV site hurts me”            “I don’t feel so good”            “I’m starting to feel itchy”            “It’s hard to breathe”            “I’m so scared. What is going on?”</p> <p><b>If learner does not stop transfusion</b>            Patient continues with dyspnea, wheezing, hypotension, discomfort with deterioration            “I feel awful; make it stop!”</p>	<p><b>Operator:</b></p> <p><b>Calls in to Sim Room:</b>            15 minutes has elapsed.</p> <p>HR: 114, weak            RR: 28            BP: 90/60            T: 99.4° F            Bilateral wheezes            O<sub>2</sub> sat: 90% on 2 L</p> <p><b>Triggers:</b>  <b>Transfusion is stopped</b></p> <p><b>If transfusion is not stopped immediately deteriorate VS:</b>            HR: 120, weak            RR: 38            B/P: 80/40            T: 99.4 F            O<sub>2</sub> sat 88% on 2L            Bilateral wheezes</p>	<p><b>Learner Actions:</b></p> <ul style="list-style-type: none"> <li>• Stays with and monitors patient for first 15 minutes of transfusion</li> <li>• Reassess VS / document</li> <li>• Recognizes transfusion reaction is occurring</li> <li>• Assess specific symptoms/ reassure patient</li> <li>• Call for help early</li> <li>• Increases oxygen to 4L</li> <li>• Notify MD immediately</li> <li>• Follows protocol for transfusion reaction               <ul style="list-style-type: none"> <li>• Stops blood transfusion</li> <li>• Prime new IV tubing with NS</li> <li>• Saves blood tubing and blood to return to blood bank per policy</li> <li>• Check information on donor blood</li> </ul> </li> </ul>	<p><b>Debriefing Points:</b></p> <ul style="list-style-type: none"> <li>• Importance of reassessment during first 10-15 minutes of transfusion</li> <li>• Significance of changes in patient status</li> <li>• Signs/symptoms of reaction/type</li> <li>• Strategies for decision making and priority setting for patients with a transfusion reaction</li> <li>• Immediate nursing interventions for patients with a blood transfusion reaction</li> <li>• Strategies for communicating with patient to decrease own and patient anxiety</li> </ul>

STATE / PATIENT STATUS	DESIRED ACTIONS & TRIGGERS TO MOVE TO NEXT STATE		
<p><b>4.</b></p> <p>Transfusion stopped; mild recovery state Patient calmer but lethargic; some chills, moaning</p>	<p><b>Operator:</b></p> <p><b>Mild recovery:</b> HR: 110 RR: 24 BP: 100/64 T: 99.2 F O<sub>2</sub> sat: 92% on 2L</p> <p><b>Triggers:</b> Notifies MD using SBAR</p>	<p><b>Learner Actions:</b></p> <ul style="list-style-type: none"> <li>• Monitor /document VS, O<sub>2</sub> sat</li> <li>• Notifies MD using SBAR</li> <li>• Follow steps of blood transfusion policy</li> <li>• Reassure patient</li> </ul>	<p><b>Debriefing Points</b></p> <ul style="list-style-type: none"> <li>• Critical factors to communicate when calling the physician using SBAR</li> <li>• Anticipate health care provider orders</li> <li>• Importance of continued assessment/monitoring</li> <li>• Evaluate effectiveness of nursing interventions</li> </ul>

Dr. provides order for:  
Epinephrine 1:1000 0.5 ml SQ (or Epi 1:10000 0.5 ml IV)  
NS bolus 500 ml over 1 hour  
Diphenhydramine 50 mg IV push x1 now

STATE / PATIENT STATUS		DESIRED ACTIONS & TRIGGERS TO MOVE TO NEXT STATE	
<p><b>5.</b> Pt states "I feel better now" "What happened?"</p>	<p><b>Operator:</b> <i>Continued recovery:</i> HR: 100 RR: 20 BP: 110/70 T: 99.2F O<sub>2</sub> Sat: 95% on 4L</p>	<ul style="list-style-type: none"> <li>• Receive MD orders</li> <li>• Read back and verifies orders to MD, accurately records new orders (or access EMR to verify new orders.)</li> <li>• Continued monitoring / assessment of VS, symptoms</li> <li>• Document on institution-specific reporting form</li> </ul> <p>If desired:</p> <ul style="list-style-type: none"> <li>• Administers medications safely using two patient identifiers and the 5 rights.</li> </ul>	<ul style="list-style-type: none"> <li>• Evaluate effectiveness of nursing interventions</li> <li>• Documentation of interventions</li> <li>• Policies/procedures for transfusion reactions</li> <li>• Policies/procedures for initiating new orders (verbal vs. EMR verification)</li> </ul> <ul style="list-style-type: none"> <li>• Safety in administering medications in a known patient</li> </ul>
<p>Scenario End Point: Learners give SBAR report of patient status to charge nurse. VS return to baseline, blood tubing returned to lab, transfusion reaction documentation completed</p>			
<p>Suggestions to <u>decrease</u> complexity: Split into 2-3 scenarios for beginning students            Suggestions to <u>increase</u> complexity: Patient can deteriorate more rapidly and have respiratory arrest. Wife can come in yelling "I told you God wouldn't want you to take someone else's blood."</p>			



**APPENDIX B: Digital images of manikin and/or scenario milieu**

<p><b>Insert digital photo here</b></p>	<p><b>Insert digital photo here</b></p>
<p><b>Insert digital photo here</b></p>	<p><b>Insert digital photo here</b></p>

## APPENDIX C: DEBRIEFING GUIDE

General Debriefing Plan			
<input type="checkbox"/> Individual	<input type="checkbox"/> Group	<input type="checkbox"/> With Video	<input type="checkbox"/> Without Video
Debriefing Materials			
<input type="checkbox"/> Debriefing Guide	<input type="checkbox"/> Objectives	<input type="checkbox"/> Debriefing Points	<input type="checkbox"/> QSEN
QSEN Competencies to consider for debriefing scenarios			
<input type="checkbox"/> Patient Centered Care	<input type="checkbox"/> Teamwork/Collaboration	<input type="checkbox"/> Evidence-based Practice	
<input type="checkbox"/> Safety	<input type="checkbox"/> Quality Improvement	<input type="checkbox"/> Informatics	
Sample Questions for Debriefing			
<ol style="list-style-type: none"> <li>1. How did the experience of caring for this patient feel for you and the team?</li> <li>2. Did you have the knowledge and skills to meet the learning objectives of the scenario?</li> <li>3. What GAPS did you identify in your own knowledge base and/or preparation for the simulation experience?</li> <li>4. What RELEVANT information was missing from the scenario that impacted your performance? How did you attempt to fill in the GAP?</li> <li>5. How would you handle the scenario differently if you could?</li> <li>6. In what ways did you check feel the need to check ACCURACY of the data you were given?</li> <li>7. In what ways did you perform well?</li> <li>8. What communication strategies did you use to validate ACCURACY of your information or decisions with your team members?</li> <li>9. What three factors were most SIGNIFICANT that you will transfer to the clinical setting?</li> <li>10. At what points in the scenario were your nursing actions specifically directed toward PREVENTION of a negative outcome?</li> <li>11. Discuss actual experiences with diverse patient populations.</li> <li>12. Discuss roles and responsibilities during a crisis.</li> <li>13. Discuss how current nursing practice continues to evolve in light of new evidence.</li> <li>14. Consider potential safety risks and how to avoid them.</li> <li>15. Discuss the nurses' role in design, implementation, and evaluation of information technologies to support patient care.</li> </ol>			
<b>Notes for future sessions:</b>			